

# **Internal Waves Over the New England Shelf**

Albert J. Plueddemann  
Department of Physical Oceanography, MS-29  
Woods Hole Oceanographic Institution  
Woods Hole, MA 02543  
Phone: (508) 289-2789 Fax: (508) 457-2181 [aplueddemann@whoi.edu](mailto:aplueddemann@whoi.edu)

Craig M. Lee  
University of Washington  
Applied Physics Laboratory  
1013 NE 40th St.  
Seattle, WA 98105-6698  
Phone: (206) 685-7656 Fax: (206) 543-6785 [craig@apl.washington.edu](mailto:craig@apl.washington.edu)

Award Number: N00014-99-1-0178 (AJP); N00014-99-1-0180 (CML)

## **LONG-TERM GOALS**

Our long-term goal is to better understand processes controlling the horizontal and vertical distribution of internal wave energy over the continental shelf. Emphasis is placed on the near-inertial band. Both the initial response to impulsive forcing and the overall distribution of near-inertial energy are of interest.

## **OBJECTIVES**

We will investigate several aspects of the internal wave field over the New England Shelf, considered to be representative of a general class of broad, gently-sloping shelves. Specifically, we intend to characterize the horizontal and vertical structure of the internal wave field over the shelf and examine the near-inertial response to impulsive wind forcing.

## **APPROACH**

Data from the Nantucket Shoals Flux Experiment (NSFE) and the combined Coastal Mixing and Optics (CMO) and Shelf Break PRIMER experiments will be used to document characteristics of near-inertial waves. Surface forcing fields will be examined to identify events that evoke strong near-inertial responses. Analytical models based on the two-layer formulations of Pettigrew (1980) and Millot and Crepon (1981) will be used as a guide to interpreting the observations. If the observed stratification warrants the additional complexity, a continuously stratified model (Kundu et. al., 1983) will be employed.

## **WORK COMPLETED**

The near-inertial signal was isolated from the CMO current meter data by removing the dominant barotropic tidal constituents and band-pass filtering the resulting record. Surface forcing events warranting further study have been identified for NSFE following Wood and Chapman (1989) and for

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 SEP 2001</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2001 to 00-00-2001</b>	
4. TITLE AND SUBTITLE <b>Internal Waves Over the New England Shelf</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Department of Physical Oceanography, MS-29,,Woods Hole Oceanographic Institution,,Woods Hole,,MA, 02543</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>Our long-term goal is to better understand processes controlling the horizontal and vertical distribution of internal wave energy over the continental shelf. Emphasis is placed on the near-inertial band. Both the initial response to impulsive forcing and the overall distribution of near-inertial energy are of interest.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>4</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			

CMO/PRIMER using the buoy meteorology and the regional model results described by Baumgartner and Anderson (1999). Aspects of surface forcing relevant to analytical modeling (front translation speeds, storm sizes, translation speeds and durations) were determined for CMO/PRIMER. A hierarchy of two-layer analytical models (of increasing complexity) has been developed and used to explore the oceanic response for a variety of model parameters. There are three models distinguished by the type of forcing: Impulsive forcing (delta function) as described by Pettigrew (1980), propagating step function forcing (representing the leading edge of a front) and propagating pulse forcing (representing the leading and trailing edges of a storm system).

## **RESULTS**

The near-inertial signals extracted from both NSFE and CMO/PRIMER show responses to surface forcing which can be approximated as a two-layer flow. There is a tendency for oscillations in the upper and lower layers to be approximately out of phase. To examine the effects of forcing by atmospheric fronts moving in the offshore direction, we used the propagating step and propagating pulse models. The response was found to be sensitive to the duration of the pulse relative to the inertial period and the speed of the front relative to the barotropic wave speed.

Initial model runs concentrated on simulating storm fronts with offshore translation (cf. Kundu and Thompson, 1985). However, for realistic translation speeds these runs always showed upper and lower layer responses which were nearly in-phase. Modifying the forcing to represent a cross-shelf uniform pulse of finite duration (by using an infinitely fast cross-shelf translation speed) showed an out of phase response more similar to the observations. This idealized forcing is consistent with wind events during CMO, which propagated rapidly alongshore, rather than offshore, and appeared at a given along-shelf location as a nearly instantaneous pulse across the entire shelf. Pulses lasting an integral number of inertial periods produce a highly damped near-inertial response due to destructive interference between waves generated by the leading and trailing edge 'fronts'. In contrast, pulses lasting half an inertial period produce constructive interference and correspondingly strong near-inertial motions.

## **IMPACT/APPLICATIONS**

By extending the analytical work done by previous investigators, we hope to elucidate the principal processes which control the near-inertial response on broad, shallow shelves. Through comparison with observations the ability of simple two-layer models to reproduce the observed response will be evaluated.

## **TRANSITIONS**

None.

## **RELATED PROJECTS**

We are using archived data from NSFE (supported by the National Marine Fisheries Service, the U.S. Geological Survey, and the National Science Foundation) and from the CMO moored array and the Shelfbreak PRIMER experiment (funded by the Office of Naval Research (ONR)). We are sharing data and results with M. Levine and T. Boyd at Oregon State University who are funded by ONR to investigate the coastal internal wave field.

## REFERENCES

- Baumgartner, M. F., and S. P. Anderson, 1999. Evaluation of NCEP regional numerical weather prediction model surface fields over the Middle Atlantic Bight. *J. Geophys. Res.*, **104**(C8), 18,141-18,158.
- Kundu, P. K., S. -Y. Chao, and J. P. McCreary, 1983. Transient coastal currents and inertio-gravity waves. *Deep-Sea Res.*, **30**, 1059-1082.
- Kundu, P. K., and R. E. Thompson, 1985. Inertial oscillations due to a moving front. *J. Phys. Oceanogr.*, **15**(8), 1076-1084.
- Millot, C., and M. Crepon, 1981. Inertial oscillations on the continental shelf of the Gulf of Lions - Observations and theory. *J. Phys. Oceanogr.*, **11**, 639-657.
- Pettigrew, N. R., 1980. The dynamics and kinematics of the coastal boundary layer off Long Island. Ph.D. Thesis, MIT/WHOI Joint Program in Oceanography, Woods Hole, MA, 262 pp.
- Wood, T. M., and D. C. Chapman, 1989. Observations of near-inertial motions during the Nantucket Shoals Flux Experiment. Unpublished manuscript, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts.